

CLAIMS

1. The use of one or more of the compounds of the group consisting of 4-thiouridine, isomaltitol, and uridine in the preparation of therapeutically effective compositions against acute or chronic inflammations, and/or problems in hemostasis related to platelet function,
5 with the exception of the use of uridine in the treatment of inflammatory conditions caused by a bacterial infection.
2. The use according to claim 1, wherein the therapeutically effective compound is 4-thiouridine.
- 10 3. The use according to claim 1, wherein the therapeutically effective compound is isomaltitol.
4. The use according to claim 1, wherein the therapeutically effective compound is uridine.
- 15 5. The use according one or more of claims 1-4, wherein the therapeutically effective dose is 1 to 100 mg per kg bodyweight.
6. Method for treatment of acute or chronic inflammations, and/or problems in hemostasis
20 related to platelet function,
characterized in that a therapeutically effective amount of one or more the of compounds of the group consisting of 4-thiouridine, isomaltitol, and uridine in the preparation of therapeutically effective compositions against acute or chronic inflammations with the
exception of the use of uridine in the treatment of inflammatory conditions caused by a
25 bacterial infection..
7. Method according to claim 6, wherein the active compound is administered in such an amount that the serum concentration thereof is 0.1 to 100 mM.

09565533.10201